

EMIS-2017 Ireland

RISK AND PRECAUTIONARY BEHAVIOURS

FINDINGS FROM THE EUROPEAN MEN WHO HAVE SEX WITH MEN
INTERNET SURVEY (IRELAND)

Read the full report at www.gayhealthnetwork.ie/research




BETWEEN **HIV** DIAGNOSIS AND
STARTING TREATMENT IN
2017



EVER TAKEN
PEP
(POST-EXPOSURE
PROPHYLAXIS)

WERE USING
PrEP
(PRE-EXPOSURE
PROPHYLAXIS)



3% DAILY


1% EVENT-BASED



48% VACCINATED AGAINST
HEPATITIS A




53% VACCINATED AGAINST
HEPATITIS B



HAD SEX WITH
**NON-STEADY
MALE PARTNER(S)**
IN THE LAST YEAR

OF THESE, **35%**
HAD CONDOMLESS
ANAL INTERCOURSE
WITH **2 OR MORE**
NON-STEADY PARTNERS



USED AT LEAST ONE
ILLICIT DRUG IN THE
LAST YEAR

CANNABIS **34%**

COCAINE **20%**

ECSTASY PILLS **19%**



USED
**STIMULANT
DRUGS**

IN THE LAST YEAR TO MAKE
SEX MORE INTENSE OR LAST
LONGER

14% OVERALL

26% OF MEN LIVING WITH HIV